



**eunethta**  
EUROPEAN NETWORK FOR HEALTH TECHNOLOGY ASSESSMENT

EUnetHTA Joint Action 3 WP4

**“Rolling Collaborative Review” of Covid-19 treatments**

**TOCILIZUMAB FOR THE TREATMENT OF COVID-19**

**Project ID: RCR03**  
Monitoring Report

**Version 8.0, March 2021**

Template version February 2021



This Rolling Collaborative Review Living Document is part of the project / joint action '724130 / EUnetHTA JA3' which has received funding from the European Union's Health Programme (2014-2020)

## DOCUMENT HISTORY AND CONTRIBUTORS

Version	Date	Description of changes
V 1.0	14/08/2020	First version
V 2.0	15/09/2020	Second version
V 3.0	15/10/2020	Third version
V 4.0	18/11/2020	Fourth version
V 5.0	15/12/2020	Fifth version
V 6.0	18/01/2021	Sixth version
V 7.0	15/02/2021	Seventh version
V 7.1	09/03/2021	Literature searches, Literature screening, Data extraction
V 7.2	10/03/2021	Data extraction and analysis complete
V 7.3	11/03/2021	Check of data extraction and analysis
V 8.0	15/03/2021	Eighth version

### Major changes from previous version

Chapter, page no.	Major changes from version 7.0
Summary	Table 4-7 Summary of safety from observational studies (AE and SAE) of tocilizumab Table 4-8 Ongoing trials of single agent tocilizumab Table 4-9 Ongoing trials of combination therapies tocilizumab

### Disclaimer

The content of this “Rolling Collaborative Review” (RCR) represents a consolidated view based on the consensus within the Authoring Team; it cannot be considered to reflect the views of the European Network for Health Technology Assessment (EUnetHTA), EUnetHTA’s participating institutions, the European Commission and/or the Consumers, Health, Agriculture and Food Executive Agency or any other body of the European Union. The European Commission and the Agency do not accept any responsibility for use that may be made of the information it contains.

### Rolling Collaborative Review team

Author(s)	National Institute of Pharmacy and Nutrition (NIPN), Hungary
Co-Author(s)	Department of Epidemiology Lazio Regional Health Service (DEPLazio), Italy

## Further contributors

Project Management	
Zorginstituut Nederland (ZIN), Netherlands	Coordination between involved parties throughout the assessment
Austrian Institute for Health Technology Assessment (AIHTA), Austria	Coordination of RCR

## Conflict of interest

All authors and co-authors involved in the production of this living document have declared they have no conflicts of interest in relation to the technology and comparator(s) assessed according to the EUnetHTA declaration of interest (DOI) form. Conflict of Interest was evaluated following the [EUnetHTA Procedure Guidance for handling DOI form \(https://eunetha.eu/doi\)](https://eunetha.eu/doi).

## Copyright

EUnetHTA assessments are published under a “CC/BY/NC” [Creative Commons Licence](https://creativecommons.org/licenses/by-nc/4.0/).



## How to cite this assessment

Please cite this assessment as follows:

EUnetHTA Rolling Collaborative (RCR03) Authoring Team. Tocilizumab for the treatment of COVID-19. Diemen (The Netherlands): EUnetHTA; 2021. 46 pages. Report No.: RCR03, Version 8.0, March 2021. Available from: [https //www.eunetha.eu](https://www.eunetha.eu).

Contact the EUnetHTA Secretariat [EUnetHTA@zinl.nl](mailto:EUnetHTA@zinl.nl) with inquiries about this assessment.

## TABLE OF CONTENTS

<b>DOCUMENT HISTORY AND CONTRIBUTORS</b> .....	<b>2</b>
<b>TABLE OF CONTENTS</b> .....	<b>4</b>
<b>LIST OF TABLES AND FIGURES</b> .....	<b>4</b>
<b>1 OBJECTIVE</b> .....	<b>6</b>
<b>2 METHODS</b> .....	<b>6</b>
2.1 SCOPE.....	6
2.2 SOURCES OF INFORMATION.....	8
<b>3 ABOUT THE TREATMENT</b> .....	<b>11</b>
3.1 MODE OF ACTION .....	11
3.2 REGULATORY STATUS .....	11
3.3 LEVEL OF EVIDENCE .....	11
<b>4 SUMMARY</b> .....	<b>12</b>
4.1 EFFECTIVENESS AND SAFETY EVIDENCE FROM RCTs.....	12
4.2 SAFETY EVIDENCE FROM OBSERVATIONAL STUDIES .....	12
4.3 ONGOING STUDIES.....	12
4.4 SCIENTIFIC CONCLUSION ABOUT STATUS OF EVIDENCE GENERATION .....	13
<b>5 REFERENCES</b> .....	<b>36</b>
<b>6 APPENDIX</b> .....	<b>39</b>
6.1 SEARCH STRATEGY TO IDENTIFY RANDOMISED CONTROLLED TRIALS.....	39
6.2 SEARCH STRATEGY TO IDENTIFY OBSERVATIONAL STUDIES.....	42
6.3 SEARCH STRATEGY TO IDENTIFY ONGOING STUDIES.....	44
6.4 FLOW DIAGRAMS .....	45

## LIST OF TABLES AND FIGURES

Table 2-1 Scope of the RCR .....	6
Table 4-1 Summary of findings (SoF) table for published RCTs related to effectiveness and safety of tocilizumab.....	14
Table 4-2. Summary of findings (SoF) table for published RCTs related to effectiveness and safety of tocilizumab.....	15
Table 4-3. Summary of findings (SoF) table for published RCTs related to effectiveness and safety of favipiravir+tocilizumab .....	15
Table 4-4. Summary of findings (SoF) table for published RCTs related to effectiveness and safety of favipiravir+tocilizumab .....	16
Table 4-5. Summary of findings (SoF) table for published RCT related to effectiveness and safety of tocilizumab.....	16
Table 6-1 Search strategy to identify randomised controlled studies.....	40
Table 6-2 Search strategy to identify observational studies.....	42
Table 6-3 Search strategy to identify ongoing studies .....	44
Appendix Figure 6-1. Flow diagram depicting the selection process of RCTs.....	45
Appendix Figure 6-2. Flow diagram depicting the selection process of observational studies.....	46

## LIST OF ABBREVIATIONS

AE	Adverse Event
ARR	Absolute Risk Reduction
ATC	Anatomical Therapeutic Chemical [Classification System]
ATMP	Advanced therapy medicinal product
CI	Confidence Interval
CRP	C-Reactive Protein
DOI	Declaration of interest
EUnetHTA	European Network of Health Technology Assessment
GRADE	Grading of Recommendations, Assessment, Development and Evaluation
HR	Hazard Ratio
HRQOL	Health-related Quality of Life
ICD	International Classification of Diseases
ICU	Intensive Care Unit
ITT	Intention-to-treat
MD	Mean Difference
MeSH	Medical Subject Headings
NA	Not applicable
NR	Not reported
OR	Odds Ratio
PP	Per Protocol
RCT	Randomized Controlled Trial
REA	Relative Effectiveness Assessment
RR	Relative Risk
SAE	Serious Adverse Event
SD	Standard Deviation
SMD	Standardized Mean Difference
SmPC	Summary of product characteristics
SoC	Standard of Care
SOP	Standard Operating Procedure
TCZ	Tocilizumab
WP4	Work Package 4

## 1 OBJECTIVE

The aim of this EUnetHTA Rolling Collaborative Review is

- to inform health policy at the national/regional and at the European level at an early stage in the life-cycle of therapies which interventions are currently undergoing clinical trials,
- to monitor (ongoing studies and their results) permanently - in the format of a Living Document - potential therapies against covid-19,
- to present comparative data on effectiveness and safety of potential therapies and
- to support preparations for an evidence-based purchasing of regional/ national health politicians, if necessary.

To avoid redundancies and duplication, the EUnetHTA Rolling Collaborative Review will reuse sources from international initiatives to collect information and data on Covid-19 treatments.

The scope of the Rolling Collaborative Review is of descriptive nature. These **EUnetHTA Rolling Collaborative Reviews are not meant to substitute a joint Relative Effectiveness Assessment (REA)** adhering to the agreed procedures and aiming at critical appraisal of the clinical evidence based on the Submission Dossier submitted by the (prospective) Marketing Authorization Holder (MAH).

## 2 METHODS

This Rolling Collaborative Review is prepared according to the project plan (“Rolling Collaborative Review (RCR) on Covid-19 treatments: Project description and planning”, published [on the EUnetHTA website](#)) and will be updated monthly. Monthly updates are published on the EUnetHTA Covid-19 Website (<https://eunethta.eu/covid-19-treatment/>) and on the EUnetHTA Rolling Collaborative Review Sharepoint page each 15<sup>th</sup> of the month.

### 2.1 Scope

**Table 2-1 Scope of the RCR**

Description	Project Scope
<p><b>Population</b></p>	<p><b>Disease</b></p> <ul style="list-style-type: none"> <li>• SARS-CoV-2 is a novel coronavirus causing a respiratory illness termed Covid-19. The full spectrum of Covid-19 ranges from mild, self-limiting respiratory tract illness to severe progressive pneumonia, multi-organ failure, and death.</li> </ul> <p><b>ICD-Codes</b> (<a href="https://www.who.int/classifications/icd/covid19/en">https://www.who.int/classifications/icd/covid19/en</a>)</p> <ul style="list-style-type: none"> <li>• An emergency ICD-10 code of ‘U07.1 COVID-19, virus identified’ is assigned to a disease diagnosis of COVID-19 confirmed by laboratory testing.</li> <li>• An emergency ICD-10 code of ‘U07.2 COVID-19, virus not identified’ is assigned to a clinical or epidemiological diagnosis of COVID-19 where laboratory confirmation is inconclusive or not available.</li> <li>• Both U07.1 and U07.2 may be used for mortality coding as cause of death. See the International guidelines for certification and classification (coding) of COVID-19 as cause of death following the link below.</li> <li>• In ICD-11, the code for the confirmed diagnosis of COVID-19 is RA01.0 and the code for the clinical diagnosis (suspected or probable) of COVID-19 is RA01.1.</li> </ul> <p><b>MeSH-terms</b></p> <ul style="list-style-type: none"> <li>• COVID-19, Coronavirus Disease 2019</li> </ul> <p><b>Target population</b> (<a href="https://www.covid19treatmentguidelines.nih.gov/overview/management-of-covid-19/">https://www.covid19treatmentguidelines.nih.gov/overview/management-of-covid-19/</a>)</p>

	<ul style="list-style-type: none"> <li>• Asymptomatic or pre-symptomatic Infection: Individuals who test positive for SARS-CoV-2 by virologic testing using a molecular diagnostic (e.g., polymerase chain reaction) or antigen test, but have no symptoms.</li> <li>• Mild Illness: Individuals who have any of the various signs and symptoms of COVID 19 (e.g., fever, cough, sore throat, malaise, headache, muscle pain) without shortness of breath, dyspnoea, or abnormal chest imaging.</li> <li>• Moderate Illness: Individuals who have evidence of lower respiratory disease by clinical assessment or imaging and a saturation of oxygen (SpO<sub>2</sub>) ≥94% on room air at sea level.</li> <li>• Severe Illness: Individuals who have respiratory frequency &gt;30 breaths per minute, SpO<sub>2</sub> &lt;94% on room air at sea level, ratio of arterial partial pressure of oxygen to fraction of inspired oxygen (PaO<sub>2</sub>/FiO<sub>2</sub>) &lt;300 mmHg, or lung infiltrates &gt;50%.</li> <li>• Critical Illness: Individuals who have respiratory failure, septic shock, and/or multiple organ dysfunction.</li> </ul>
<p><b>Intervention</b></p>	<p>Tocilizumab is a humanised IgG1 monoclonal antibody that binds specifically to both soluble and membrane-bound IL-6 receptors (sIL-6R and mL-6R) and inhibits sIL-6R and mL-6R-mediated signalling. Tocilizumab is indicated (EMA-approved) for the treatment of</p> <ul style="list-style-type: none"> <li>• rheumatoid arthritis in adults</li> <li>• giant cell arteritis in adults</li> <li>• active systemic juvenile idiopathic arthritis in patients aged ≥2 years</li> <li>• juvenile idiopathic polyarthritis in patients aged ≥2 years</li> </ul> <p>chimeric antigen receptor (CAR) T cell-induced severe or life-threatening cytokine release syndrome (CRS) in patients aged ≥2 years</p>
<p><b>Comparison</b></p>	<p>Any active treatment, placebo, or standard of care.</p> <p><b>Rationale:</b> Since there is no gold standard treatment any comparator is acceptable as well as the above listed interventions.</p>
<p><b>Outcomes</b></p>	<p><u>Main outcome:</u></p> <ul style="list-style-type: none"> <li>• All-cause Mortality (Survival)</li> </ul> <p><u>Additional Outcomes:</u></p> <p>Efficacy:</p> <ul style="list-style-type: none"> <li>• Length of hospital stay,</li> <li>• Viral burden (2019-nCoV RT-PCR negativity),</li> <li>• Clinical progression (WHO Clinical Progression Scale measured daily over the course of the study),</li> <li>• Rates of hospitalization and of patients entering ICU,</li> <li>• Duration of mechanical ventilation,</li> <li>• Quality of life.</li> </ul> <p>Safety:</p> <ul style="list-style-type: none"> <li>• Adverse events (AE),</li> <li>• Severe adverse events (SAE),</li> <li>• Withdrawals due to AEs,</li> <li>• Most frequent AEs,</li> <li>• Most frequent SAEs.</li> </ul> <p><b>Rationale:</b> We will give priority according to the Core Outcome Set for Clinical Trials on Coronavirus Disease 2019 (<a href="https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7102592/pdf/main.pdf">https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7102592/pdf/main.pdf</a>) and A minimal common outcome measure set for COVID-19 clinical research from the WHO Working Group on the Clinical Characterisation and Management of COVID-19 infection.</p>

<b>Study design</b>	Efficacy: randomised controlled trials (RCT) Safety: observational studies (comparative or single-arm prospective studies and registries)
---------------------	--

## 2.2 Sources of information

According to the project plan, this Rolling Collaborative Review is based on three main sources of information, as described below:

### 1. Summary of findings(SoF) table for published RCTs related to effectiveness and safety:

This table is based on the living systematic review and Network Meta-Analysis (NMA) created by the partnering institute of DEPLazio: [find the PROSPERO protocol here](#). DEPLazio provides updates for the SoF table on a monthly basis to the EUnetHTA partners authoring the respective Rolling CR documents who are integrating this information accordingly.

The literature search is conducted in the following databases:

- PubMed
- MEDLINE, accessed via OVID
- Embase, accessed via OVID

<b>Population</b>	People affected by COVID-19, as defined by the authors of the studies. No limits in terms of gender or ethnicity.  SARS-CoV-2 is a novel coronavirus causing a respiratory illness termed Covid-19. It started spreading in December 2019, and was declared a pandemic by the World Health Organisation on 11th March 2020. The full spectrum of Covid-19 ranges from mild, self-limiting respiratory tract illness to severe progressive pneumonia, multi-organ failure, and death.
<b>Intervention</b>	Interventions for the treatment of people affected by COVID-19, including pharmacological interventions (e.g. antibiotics, antibodies, antimalarial, antiviral, antiretroviral, immune-suppressors/modulators, kinase inhibitors) and their combinations.
<b>Comparison</b>	Any active treatment, placebo, or standard of care.
<b>Outcomes</b>	All-cause mortality  Additional outcomes: Length of hospital stay, 2019-nCoV RT-PCR negativity, PaO <sub>2</sub> /FiO <sub>2</sub> , Duration of mechanical ventilation, radiological imaging, Adverse events, Severe adverse events.
<b>Study design</b>	Randomised controlled trials (RCT); no restriction on language of publication

To identify preprints of preliminary reports of work that have not been peer-reviewed, the following sources are searched:

- medRxiv Health Sciences
- bioRxiv Biology

In addition to the sources and strategies described above, registers of ongoing studies are screened. Key conferences and conference proceedings are considered. Appendix Table 6-1 describes in detail the sources searched, the search terms used and the dates at which the searches are executed.



Data extraction, Risk of bias assessment, data synthesis:

Two reviewers from DEPLazio are screening search results, assessing full texts of studies and extract study characteristics and outcome data according to pre-defined criteria. The process of study selection is depicted as a flow diagram in Appendix Figure 6-1.

Risk of bias is assessed using the criteria outlined in the Cochrane Handbook for Systematic Reviews of Interventions [1].

Dichotomous outcomes are analysed by calculating the relative risk (RR) for each trial with the uncertainty in each result being expressed by its 95% confidence interval (CI). Continuous outcomes are analysed by calculating the mean difference (MD) with the relative 95% CI when the study used the same instruments for assessing the outcome.

The standardised mean difference (SMD) is applied when studies used different instruments. Pairwise meta-analyses is performed for primary and secondary outcomes using a random-effects model in RevMan for every treatment comparison [2]. Network meta-analysis (NMA) is performed for the primary outcome. For rating the certainty of the evidence, the GRADE approach is being used [3].

- Sources: <http://deplazio.net/farmacicovid/index.html> for SoF (or <https://covid-nma.com/>)

**2. Table(s) on published (peer reviewed) observational studies for safety results:**

The literature search is conducted on a monthly basis.

The sources and search methods are described in more detail in Appendix Table 6-2.

<b>Population</b>	See project Scope
<b>Intervention</b>	Tocilizumab is a humanised IgG1 monoclonal antibody that binds specifically to both soluble and membrane-bound IL-6 receptors (sIL-6R and mIL-6R) and inhibits sIL-6R and mIL-6R-mediated signaling.
<b>Comparison</b>	Any active treatment, placebo, or standard of care.
<b>Outcomes</b>	See project Scope
<b>Study design</b>	Inclusion criteria: Prospective non-randomised controlled trials, prospective case series (i.e. comparative or single-arm prospective studies), registries  Exclusion criteria: retrospective studies, case studies/ case reports, observational studies that do not report safety data

Two researchers from NIPHNO carry out title and abstract screening and assess the full texts of all potentially eligible studies. The study selection process is depicted in a flow diagram (Appendix Figure 6-2).

One researcher of NIPN extracts the data and assesses the risk of bias using Robins-I (<https://training.cochrane.org/handbook/current/chapter-25>). For prospective single arm studies the Newcastle-Ottawa Scale (NOS) is used to assess the methodological rigor and applicability.

Results are presented in tabular form for all included studies.

**3. Table(s) on ongoing trials:**

The following clinical trial registries are searched on a monthly basis:

- ClinicalTrials.gov: <https://clinicaltrials.gov/>
- ISRCTN: <https://www.isrctn.com/>
- European Clinical Trials Registry: <https://www.clinicaltrialsregister.eu/>

Inclusion criteria: Randomised controlled trials, Controlled trials

One researcher of NIPN is searching and extracting the data for the eligible studies.. At the drafting stage of each update, the author team verifies whether the status of previously identified studies has changed. This is done by verifying the date of the last update posted in the trial registers. In addition, trial register IDs of all previously identified studies are entered in both PubMed and Google (google.com) to verify if previously identified studies have been published since the last update. In Google, the first 10 hits are screened for this purpose.

Search methods are described in more detail in Table 6-3.

Data are presented in tabular form.

## 3 ABOUT THE TREATMENT

### 3.1 Mode of Action

Tocilizumab is a humanised IgG1 monoclonal antibody that binds specifically to both soluble and membrane-bound IL-6 receptors (sIL-6R and mIL-6R) and inhibits sIL-6R and mIL-6R-mediated signaling. IL-6 is a pleiotropic, pro-inflammatory cytokine produced by a variety of cell types, including lymphocytes, monocytes, and fibroblasts. Infection by the related SARS-associated coronavirus induces a dose-dependent production of IL-6 from bronchial epithelial cells. Elevations in IL-6 levels may be an important mediator when severe systemic inflammatory responses occur in patients with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. COVID-19-associated systemic inflammation and hypoxic respiratory failure is associated with heightened cytokine release, as indicated by elevated blood levels of IL-6, C-reactive protein (CRP), D-dimer, and ferritin [4].

### 3.2 Regulatory Status

The Market Authorisation Holder of tocilizumab is Roche Pharma. Tocilizumab is not approved by the European Medicines Agency (EMA) or the American Food and Drug Administration (FDA) for COVID-19 patients. Tocilizumab is indicated (EMA-approved) for the treatment of:

- rheumatoid arthritis in adults
- giant cell arteritis in adults
- active systemic juvenile idiopathic arthritis in patients aged  $\geq 2$  years
- juvenile idiopathic polyarthritis in patients aged  $\geq 2$  years
- chimeric antigen receptor (CAR) T cell-induced severe or life-threatening cytokine release syndrome (CRS) in patients aged  $\geq 2$  years [5].

Tocilizumab is not authorised in Covid-19 patients (EMA, FDA).

### 3.3 Level of Evidence

Sixty-three hospitalized adult patients with COVID-19 were enrolled in a prospective, open-label study of tocilizumab for severe COVID-19. Patients received either TCZ IV (8 mg/kg) or SC (324 mg); (the optional second dose within 24 hours 52 of 63 patients), and all of the patients received off-label antiretroviral protease inhibitors. Following administration of tocilizumab, fevers resolved in all but one patient, and CRP, ferritin, and D-dimer levels declined. The mean PaO<sub>2</sub>/FiO<sub>2</sub> ratio of the patients increased between admission (152 +/- 53 mm Hg) and Day 7 of hospitalization (284 +/- 116 mm Hg). No moderate or severe adverse events attributable to tocilizumab were reported. The overall mortality rate was 11% (7 of 63 patients). No details were provided regarding the rate of secondary infections after tocilizumab use. The authors report an association between earlier use of tocilizumab and reduced mortality; however, interpretation of this result is limited because the study results did not describe a comparison group or specify an a priori comparison [6].

The phase III COVACTA (NCT04320615) study of tocilizumab did not meet its primary endpoint of improved clinical status in hospitalised adult patients with severe COVID-19 associated pneumonia. In addition, the key secondary endpoints, which included the difference in patient mortality at week four, were not met; however, there was a positive trend in time to hospital discharge in patients treated with tocilizumab. The COVACTA study did not identify any new safety signals for tocilizumab.

The phase III REMDACTA (NCT04409262) is currently ongoing, and participants are receiving treatments. The phase III EMPACTA (NCT04372186) study has interim results available, and participants are receiving an intervention or are being examined, but potential participants are not currently being recruited or enrolled. The EMPACTA study showed that patients with COVID-19

associated pneumonia who received tocilizumab plus standard of care were less likely to progress to mechanical ventilation or death compared to patients who received placebo plus standard of care (log-rank p-value = 0.0348; HR [95% CI] = 0.56 [0.32, 0.97]). The cumulative proportion of patients who progressed to mechanical ventilation or death by day 28 was 12.2% in the tocilizumab arm versus 19.3% in the placebo arm. The EMPACTA study did not identify any new safety signals for tocilizumab [28].

A phase III trial involving 243 patients who require hospital but not mechanical ventilation evaluates the effects of tocilizumab compared to placebo. Tocilizumab was not effective for preventing intubation or death, the hazard ratio for intubation or death in the tocilizumab group as compared with the placebo group was 0.83 (95% confidence interval [CI], 0.38 to 1.81; P=0.64) (HR 0.83;95% confidence interval [CI], 0.38 to 1.81; P=0.64) [11]. The TOBICRAS phase III randomized clinical trial was terminated early, after 129 patients had been enrolled, because of an increased number of deaths at 15 days in the tocilizumab group [12].

The RECOVERY trial (NCT04381936) is an ongoing randomised trial investigating whether potential treatments for COVID-19 reduce the risk of death. Participants with progressive COVID-19 may undergo an optional subsequent randomisation between tocilizumab and no additional treatment. A total of 2022 patients were randomly allocated to receive tocilizumab by intravenous infusion and were compared with 2094 patients randomly allocated to usual care alone. 82% of patients were taking a systemic steroid such as dexamethasone. Treatment with tocilizumab significantly reduced deaths: 596 (29%) of the patients in the tocilizumab group died within 28 days compared with 694 (33%) patients in the usual care group (rate ratio 0.86; [95% confidence interval [CI] 0.77 to 0.96]; p=0.007), an absolute difference of 4%. The study also showed that tocilizumab shortens the time until patients are successfully discharged from hospital and reduces the need for a mechanical ventilator ([https://www.recoverytrial.net/files/recovery-press-release-tocilizumab\\_final.pdf](https://www.recoverytrial.net/files/recovery-press-release-tocilizumab_final.pdf)).

One ongoing RCT evaluates anakinra alone and in combination with siltuximab or tocilizumab.

## 4 SUMMARY

### 4.1 Effectiveness and Safety evidence from RCTs

There are insufficient data from clinical trials on the use of tocilizumab in patients with COVID-19. The data currently available are presented in Table 4-1, Table 4-2, Table 4-3, Table 4-4 and Table 4-5. The currently available evidence on all-cause mortality, frequency of adverse events, duration of hospitalization, disease severity and hospital discharges is not conclusive, as there is no statistically significant association between these outcomes and the treatment with tocilizumab. For mortality, any adverse events and hospitalization, studies show a trend towards favouring standard of care over tocilizumab, whereas for severe adverse events and the progression of disease, study results tend to favour tocilizumab.

### 4.2 Safety evidence from observational studies

The primary laboratory abnormalities reported with tocilizumab treatment are elevated liver enzyme levels that appear to be dose dependent. Neutropenia or thrombocytopenia are uncommon. Additional AEs, such as risk for serious infections (e.g., TB, other bacterial pathogens), have been reported only in the context of continuous dosing of tocilizumab. In two prospective cohort studies with high risk of bias, safety evidence has been reported. In a retrospective analysis of data from 21 patients, no adverse reaction were observed during the treatment [7]. During the 10-day follow-up Toniati et al. 2020 recorded three cases of severe adverse events: two patients developed septic shock and died, one had gastrointestinal perforation requiring urgent surgery and was alive at day 10 [8].

### 4.3 Ongoing studies

Several RCTs and interventional nRCTs related to tocilizumab alone or in combination therapy are currently ongoing.

#### **4.4 Scientific conclusion about status of evidence generation**

High quality evidence from ongoing RCTs are expected to assess effectiveness and safety of tocilizumab in COVID-19 patients.

Future controlled trials in patients with severe illness are needed to confirm or exclude the possibility of treatment benefit with tocilizumab.

**Table 4-1 Summary of findings (SoF) table for published RCTs related to effectiveness and safety of tocilizumab**

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of evidence	Comments
	Risk with standard of care	Risk with tocilizumab				
All-cause mortality [9]; [10]; [11]; [12]; [13]; [14]; [15]	209 per 1000	213 per 1000 (163 to 200)	RR 1.02 (0.78 to 1.34)	2200 (7 RCTs)	moderate	Compared to SoC there is no effect on all cause mortality
Number of patients with any adverse event [9]; [12]; [13]; [14]; [16]	557 per 1000	591 per 1,000 (466 to 746)	RR 1.06 (0.80 to 1.34)	1153 (5 RCTs)	very low	Compared to standard treatment could increase the number of patients with adverse events
Number of patients with severe adverse events [9]; [10]; [11]; [12]; [13]; [14]; [15]; [16]	144 per 1000	130 per 1,000 (108 to 156)	RR 0.90 (0.75 to 1.08)	2274 (8 RCTs)	moderate	Compared to SoC tocilizumab probably reduces the risk of severe adverse events
Length of stay in hospital [9]; [13]; [15]	Rosas 2020: HR 1.35 [95% CI (1.02; 1.79) Salama 2020: HR 1.16 [95% CI (0.91; 1.48)] REMAP study: HR 1.41 [95% CI (1.18; 1.68)] Cumulatively length of hospitalization differs between the two groups in favour of Tocilizumab (HR: 1.32 (95% CI 1.16, 1.49) p <0.0001)			(3 RCTs)	moderate	Compared to SoC there is no effect on the number of days of hospitalization
Length of stay in hospital (mean days) [12]	SMD 0.42 lower (0.77 lower to 0.07 lower)			129 (1 RCT)	low	Compared to SoC there is no effect on the number of days of hospitalization
Progression of disease severity [10]; [11]; [14]	161 per 1,000	110 per 1,000 (69 to 176)	RR 0.68 (0.43 to 1.09)	495 (3 RCTs)	moderate	Compared to SoC tocilizumab probably reduces the risk of progression of disease severity
Number of patients discharged [10]; [11]	889 per 1,000	898 per 1,000 (836 to 969)	RR 1.01 (0.94 to 1.09)	365 (2 RCTs)	moderate	Compared to SoC there is no effect on the number of patients discharged
Duration of hospitalization in intensive care [15]	HR 1.42 [IC95% (1.18; 1.71) in favor of tocilizumab			( RCT)1	moderate	Compared to SoC tocilizumab probably reduces the number of days of hospitalization in intensive care

**Source:** Cruciani F, De Crescenzo F, Vecchi S, Saulle R, Mitrova Z, Amato L, Davoli M

**Abbreviations:** CI=Confidence interval; RR=Risk ratio

**Table 4-2. Summary of findings (SoF) table for published RCTs related to effectiveness and safety of tocilizumab**

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of evidence	Comments
	Risk with favipiravir	Risk with tocilizumab				
Number of patients with any adverse event [17]	284 per 1000	400 per 1000	RR 0.71 (0.15 to 3.50)	12 (1 RCT)	very low	
Number of patients with serious adverse events [17]	No serious adverse event reported				very low	
Number of patients with significant improvement in lung disease on CT [17]	The study reports that there was a significant difference in favour of favipiravir (P = 0.034), HR 3.16 [95% CI 0.62-16.10]				very low	

**Source:** Cruciani F, De Crescenzo F, Vecchi S, Saulle R, Mitrova Z, Amato L, Davoli M

**Abbreviations:** CI=Confidence interval; RR=Risk ratio

**Table 4-3. Summary of findings (SoF) table for published RCTs related to effectiveness and safety of favipiravir+tocilizumab**

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of evidence	Comments
	Risk with favipiravir	Risk with favipiravir+tocilizumab				
Number of patients with any adverse event [17]	286 per 1000	643 per 1000	RR 2.25 (0.65 to 7.73)	21 (1 RCT)	very low	
Number of patients with serious adverse events [17]	No serious adverse event reported				very low	
Number of patients with significant improvement in lung disease on CT [17]	The study reports that the cumulative rate of lung lesion remission on day 14 was significantly higher in the combined group than in the favipiravir group (HR 2.66 95% CI [1.08-6.53], P = 0.019).				very low	

**Source:** Cruciani F, De Crescenzo F, Vecchi S, Saulle R, Mitrova Z, Amato L, Davoli M

**Abbreviations:** CI=Confidence interval; RR=Risk ratio

**Table 4-4. Summary of findings (SoF) table for published RCTs related to effectiveness and safety of favipiravir+tocilizumab**

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of evidence	Comments
	Risk with tocilizumab	Risk with favipiravir+tocilizumab				
Number of patients with any adverse event [17]	400 per 1000	644 per 1000	RR 1.61 (0.51 to 5.04)	19 (1 RCT)	moderate	
Number of patients with serious adverse events [17]	No serious adverse event reported				very low	
Number of patients with significant improvement in lung disease on CT [17]	There was no significant difference in the cumulative remission rate of lung lesions on day 14 between the combination group and the tocilizumab group HR 1.28 [95% CI 0.39-4.23] P = 0.575				very low	

**Source:** Cruciani F, De Crescenzo F, Vecchi S, Saulle R, Mitrova Z, Amato L, Davoli M

**Abbreviations:** CI=Confidence interval; RR=Risk ratio

**Table 4-5. Summary of findings (SoF) table for published RCT related to effectiveness and safety of tocilizumab**

Outcome	Anticipated absolute effects (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of evidence	Comments
	Risk with sarilumab	Risk with tocilizumab				
All-cause mortality [15]	222 per 1000	280 per 1000 (158 to 496)	RR 1.26 (0.71 to 2.23) 1	395 (1 RCT)	moderate	Compared to sarilumab tocilizumab probably increase the risk of mortality
Number of patients with serious adverse events [15]	0 per 1.000	0 per 1000 (0 to 0)	RR 2.63 (0.16 to 44.48)	401 (1 RCT)	low	Compared to sarilumab tocilizumab probably increase the risk of serious adverse events

**Source:** Cruciani F, De Crescenzo F, Vecchi S, Saulle R, Mitrova Z, Amato L, Davoli M

**Abbreviations:** CI=Confidence interval; RR=Risk ratio



Certainty assessment							N <sup>o</sup> of patients		Effect		Certainty
N <sup>o</sup> of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Tocilizumab	Standard treatment	Relative (95% CI)	Absolute (95% CI)	
<b>All-cause mortality</b>											
7 1,2,3,4,5,6,7	randomised trials	serious <sup>a</sup>	not serious	not serious	not serious	none	214/1249 (17.1%)	199/951 (20.9%)	<b>RR 1.02</b> (0.78 to 1.34)	<b>4 more per 1.000</b> (from 46 fewer to 71 more)	⊕⊕⊕○ MODERATE
<b>Number of patients with any adverse event</b>											
5 <sup>1,4,5,7,8</sup>	randomised trials	serious <sup>b</sup>	very serious <sup>c</sup>	not serious	not serious	none	432/715 (60.4%)	244/438 (55.7%)	<b>RR 1.06</b> (0.84 to 1.34)	<b>33 more per 1.000</b> (from 89 fewer to 189 more)	⊕○○○ VERY LOW
<b>Number of patients with severe adverse events</b>											
8 1,2,3,4,5,6,7,8	randomised trials	serious <sup>d</sup>	not serious	not serious	not serious	none	210/1289 (16.3%)	142/985 (14.4%)	<b>RR 0.90</b> (0.75 to 1.08)	<b>14 fewer per 1.000</b> (from 36 fewer to 12 more)	⊕⊕⊕○ MODERATE
<b>Length of stay in hospital</b>											
3 <sup>1,4,6</sup>	randomised trials	serious <sup>e</sup>	not serious	not serious	not serious	none	Rosas 2020: HR 1.35 [95% CI (1.02; 1.79)] Salama 2020: HR 1.16 [95% CI (0.91; 1.48)] REMAP study: HR 1.41 [95% CI (1.18; 1.68)] Cumulatively length of hospitalization differs between the two groups in favour of Tocilizumab (HR: 1.32 (95% CI 1.16, 1.49) p <0.0001)				⊕⊕⊕○ MODERATE

Certainty assessment							№ of patients		Effect		Certainty
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Tocilizumab	Standard treatment	Relative (95% CI)	Absolute (95% CI)	

#### Length of stay in hospital (mean days)

1 <sup>7</sup>	randomised trials	serious <sup>f</sup>	not serious	not serious	serious <sup>g</sup>	none	65	64		<b>SMD 0.42 lower</b> (0.77 lower to 0.07 lower)	⊕⊕○○ LOW
----------------	-------------------	----------------------	-------------	-------------	----------------------	------	----	----	--	---	-------------

#### Progression of COVID-19 disease

3 <sup>2,3,5</sup>	randomised trials	serious <sup>h</sup>	not serious	not serious	not serious	none	28/284 (9.9%)	34/211 (16.1%)	<b>RR 0.68</b> (0.43 to 1.09)	<b>52 fewer per 1.000</b> (from 92 fewer to 15 more)	⊕⊕⊕○ MODERATE
--------------------	-------------------	----------------------	-------------	-------------	-------------	------	---------------	----------------	----------------------------------	---	------------------

#### Number of patients discharged

2 <sup>2,3</sup>	randomised trials	serious <sup>i</sup>	not serious	not serious	not serious	none	199/221 (90.0%)	128/144 (88.9%)	<b>RR 1.01</b> (0.94 to 1.09)	<b>9 more per 1.000</b> (from 53 fewer to 80 more)	⊕⊕⊕○ MODERATE
------------------	-------------------	----------------------	-------------	-------------	-------------	------	-----------------	-----------------	----------------------------------	---	------------------

#### Duration of hospitalization in intensive care

1 <sup>6</sup>	randomised trials	serious <sup>j</sup>	not serious	not serious	not serious	none	HR 1.42 [IC95% (1.18; 1.71) in favor of tocilizumab]			⊕⊕⊕○ MODERATE
----------------	-------------------	----------------------	-------------	-------------	-------------	------	--	--	--	------------------

#### Explanations

- a. Downgraded of one level for performance bias at high risk in four studies and at unclear risk in two studies, unclear risk of selection bias in three studies, unclear risk of detection bias in two studies

- b. Downgraded of one level for performance bias at high risk in three studies and at unclear risk in one study, unclear selection bias in three studies, unclear detection bias in two studies and unclear attrition and reporting bias in another study
- c. Downgraded of two levels for high heterogeneity:  $i^2=69\%$
- d. Downgraded of one level for performance bias at high risk in five studies and at unclear risk in two studies, unclear selection bias in four studies, unclear detection bias in two studies and unclear attrition and reporting bias in another study
- e. Downgraded of one level for performance bias at high risk in one study and unclear in another, unclear risk of selection bias in two studies, unclear risk of detection bias in one study
- f. Downgraded of one level for performance bias at high risk and at unclear risk of detection bias
- g. Downgraded of one level for small sample size (<200)
- h. Downgraded of one level for performance bias at high risk in two studies and unclear in one study
- i. Downgraded of one level for performance bias at high risk in one study and unclear in another
- j. Downgraded of one level for performance bias at high risk

Source: [15]

Certainty assessment							№ of patients		Effect		Certainty
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Favipiravir	Tocilizumab	Relative (95% CI)	Absolute (95% CI)	

**Number of patients with any adverse event**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	2/7 (28.6%)	2/5 (40.0%)	<b>RR 0.71</b> (0.15 to 3.50)	<b>116 fewer per 1.000</b> (from 340 fewer to 1.000 more)	⊕○○○ VERY LOW
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	-------------	-------------	----------------------------------	--	------------------

**Number of patients with serious adverse events**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	No serious adverse event reported			⊕○○○ VERY LOW
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	-----------------------------------	--	--	------------------

**Number of patients with significant improvement in lung disease on CT**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	The study reports that there was a significant difference in favour of favipiravir (P = 0.034), HR 3.16 [95% CI 0.62-16.10]			⊕○○○ VERY LOW
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	---	--	--	------------------

**Explanations**

a. Downgraded of two levels for high risk of performance bias and unclear risk of selection bias

b. Downgraded of two levels for very small sample size

Source: [17]

Certainty assessment							No of patients		Effect		Certainty	Importance
No of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Favipiravir+Tocilizumab	Favipiravir	Relative (95% CI)	Absolute (95% CI)		

**Number of patients with any adverse event**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	9/14 (64.3%)	2/7 (28.6%)	RR 2.25 (0.65 to 7.73)	357 more per 1.000 (from 100 fewer to 1.000 more)	⊕○○○ VERY LOW	
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	--------------	-------------	------------------------	---	------------------	--

**Number of patients with serious adverse events**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	No serious adverse event reported			⊕○○○ VERY LOW	
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	-----------------------------------	--	--	------------------	--

**Number of patients with significant improvement in lung disease on CT**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	The study reports that the cumulative rate of lung lesion remission on day 14 was significantly higher in the combined group than in the favipiravir group (HR 2.66 95% CI [1.08-6.53], P = 0.019).			⊕○○○ VERY LOW	
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	---	--	--	------------------	--

**Explanations**

a. Downgraded of two levels for high risk of performance bias and unclear risk of selection bias

b. Downgraded of two levels for very small sample size

Source: [17]

Certainty assessment							№ of patients		Effect		Certainty
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Favipiravir+Tocilizumab	Tocilizumab	Relative (95% CI)	Absolute (95% CI)	

**Number of patients with any adverse event**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	9/14 (64.3%)	2/5 (40.0%)	RR 1.61 (0.51 to 5.04)	<b>244 more per 1.000</b> (from 196 fewer to 1.000 more)	⊕○○○ VERY LOW
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	--------------	-------------	------------------------	--	------------------

**Number of patients with serious adverse events**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	No serious adverse event reported				⊕○○○ VERY LOW
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	-----------------------------------	--	--	--	------------------

**Number of patients with significant improvement in lung disease on CT**

1 <sup>1</sup>	randomised trials	very serious <sup>a</sup>	not serious	not serious	very serious <sup>b</sup>	none	There was no significant difference in the cumulative remission rate of lung lesions on day 14 between the combination group and the tocilizumab group HR 1.28 [95% CI 0.39-4.23] P = 0.575				⊕○○○ VERY LOW
----------------	-------------------	---------------------------	-------------	-------------	---------------------------	------	---	--	--	--	------------------

**Explanations**

a. Downgraded of two levels for high risk of performance bias and unclear risk of selection bias

b. Downgraded of two levels for very small sample size

Source [16]

Certainty assessment							№ of patients		Effect		Certainty
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Tocilizumab	Sarilumab	Relative (95% CI)	Absolute (95% CI)	

**All-cause mortality**

1 <sup>1</sup>	randomised trials	serious <sup>a</sup>	not serious	not serious	not serious	none	98/350 (28.0%)	10/45 (22.2%)	RR 1.26 (0.71 to 2.23)	<b>58 more per 1.000</b> (from 64 fewer to 273 more)	⊕⊕⊕○ MODERATE
----------------	-------------------	----------------------	-------------	-------------	-------------	------	----------------	---------------	------------------------	--	------------------

Certainty assessment							№ of patients		Effect		Certainty
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Tocilizumab	Sarilumab	Relative (95% CI)	Absolute (95% CI)	
<b>Number of patients with serious adverse events</b>											
1 <sup>1</sup>	randomised trials	serious <sup>a</sup>	not serious	not serious	serious <sup>b</sup>	none	9/353 (2.5%)	0/48 (0.0%)	<b>RR 2.63</b> (0.16 to 44.48)	<b>0 fewer per 1.000</b> (from 0 fewer to 0 fewer)	⊕⊕○○ LOW

**Explanations**

- a. Downgraded of one level for high risk of performance bias
- b. Downgraded of one level for wide CI

**Source:** [15]

**Table 4-6 Study characteristics of included RCTs**

<b>Author, year, reference number/Study name/Study ID</b>	Rosas et al 2020 [9] COVACTA NCT04320615	Salvarani et al 2020 [10] NCT04346355	Stone et al 2020 [11] NCT04356937	Salama et al 2020 [13] EMPACTA NCT04372186	Hermine et al 2020 [14] NCT04331808	Gordon et al 2021 [15] REMAP-CAP	Veiga et al. 2021 [12] TOCIBRAS	Wang et al 2020 [16] ChiCTR2000029765	Zhao et al. 2020 [17] NCT04310228 ChiCTR2000030894
<b>Study design, study phase</b>	randomized, double-blind, placebo-controlled, phase 3	open-label, randomized clinical trial, phase 2	randomized, double-blind, placebo-controlled phase 3 trial	randomized, double-blind, placebo-controlled phase 3 trial	cohort-embedded, investigator-initiated, open-label, bayesian randomized phase 2 clinical trial	phase IV randomized, open label; multifactorial, adaptive platform trial	prospective, randomized, superiority, open-label, controlled trial	randomized controlled phase 4 trial	single-arm, open-label, phase 2
<b>Centres (single centre or multicentre), country, setting</b>	global, multicenter	multicenter, Italy	multicenter, United States	global, multicenter	multicenter, France	international	multicenter, Brazil	multicenter, China	multicentre China
<b>Patient population (number of included patients/ Mean age and sex/ Disease severity*)</b>	452 patients <b>TCZ:</b> age mean (SD) 60.9 (14.6) 69.7% male <b>placebo:</b> age mean (SD) 60.6 (13.7); 70.1% male; severe	126 patients median (range) age of 60.0 (53.0-72.0) years 61.1% male	243 patients; median age 59.8 years (range, 21.7 to 85.4); 58% male	389 patients TCZ: 60.2% male; mean ( $\pm$ SD) age 56.0 $\pm$ 14.3 years placebo:57% male; mean ( $\pm$ SD) age 55.6 $\pm$ 14.9 years	131 patients median (IQR) age was 64 (57.1-74.3) years 68% male moderate, severe	353 (toc vs, sarilumab) patients critically ill COVID-19. mean age 61,5 (SD: 12,5) years; 73,9% male	129 patients mean age 57 (SD 14) years; 68% men	65 patients	26 patients; median 73.5 years (34–89) 53.8 % (14/26) male Common, severe or critical
<b>Inclusion criteria</b>	18 years or older with severe COVID-19 pneumonia confirmed by positive polymerase	18 years and older, with COVID-19 pneumonia confirmed by positive polymerase chain reaction test;	19 to 85 years of age with COVID-19 pneumonia confirmed by positive polymerase chain	18 years and older, with COVID-19 pneumonia confirmed by positive polymerase chain reaction test	18 years and older, with COVID-19 pneumonia confirmed by positive polymerase chain reaction test	Critically ill, >18 years, with Covid-19, admitted to an intensive care unit (ICU) and receiving	severe or critically ill need for oxygen supplementation on less than 24 hours before the randomization	18 to 85 years; diagnosed with the common type of NCP (including severe risk factors) and severe cases of new coronavirus pneumonia	Laboratory-confirmed cases according to Chinese guidelines of COVID-19; Male or female more than 18 years old; Increased interleukin-6;Sign

<b>Author, year, reference number/Study name/Study ID</b>	Rosas et al 2020 [9] COVACTA NCT04320615	Salvarani et al 2020 [10] NCT04346355	Stone et al 2020 [11] NCT04356937	Salama et al 2020 [13] EMPACTA NCT04372186	Hermine et al 2020 [14] NCT04331808	Gordon et al 2021 [15] REMAP-CAP	Veiga et al. 2021 [12] TOCIBRAS	Wang et al 2020 [16] ChiCTR2000029765	Zhao et al. 2020 [17] NCT04310228 ChiCTR2000030894
	chain reaction test	acute respiratory distress syndrome	reaction test; requiring hospital but not mechanical ventilation			respiratory or cardiovascular organ support	positive inflammatory tests		the informed consent
<b>Exclusion criteria</b>	progression to death is imminent and inevitable within the next 24 hours; active tuberculosis or bacterial, fungal, or viral infection	ICU admission, known hypersensitivity to tocilizumab, and any condition preventing future admission to ICU, such as advanced age with multiple comorbidities, as well as the patient's expressed will to avoid future intubation	Uncontrolled bacterial, fungal, or non-COVID viral infection, active TB	progression to death is imminent and inevitable within the next 24 hours; active tuberculosis or bacterial, fungal, or viral infection	acute kidney injury, cardiovascular condition, pulmonary disease, obesity, high blood pressure, diabetes, chronic kidney diseases, hematological diseases, sickle cell diseases, autoimmune and auto-inflammatory, pregnant women, HIV infected	presumption that death was imminent with lack of commitment to full support, and prior participation in REMAP-CAP within 90 days.	need for mechanical ventilation for 24 hours or more before the randomization; active uncontrolled infection; liver disease, cirrhosis or elevated AST or ALT above 5 times the upper level limit; renal disease with estimate glomerular filtration below 30 mL/min/1.72 m <sup>2</sup>	pregnant or lactating women; 3. ALT / AST > 5 ULN, neutrophils < 0.5, platelets less than 50; diagnosis of rheumatic immune-related diseases; active pulmonary tuberculosis, with definite bacterial and fungal infections.	Allergic to favipiravir or tocilizumab; Pregnant or lactating woman; ALT or AST > 5 times of upper limit of normal; Patients with active hepatitis, tuberculosis, and definite bacterial or fungal infections; Other conditions judged by the investigators.
<b>Intervention (generic drug name and dosage, time frame;</b>	tocilizumab infusion of TCZ, dosed at 8 mg/kg, up to a	TCZ: 8 mg/kg up to a maximum of 800 mg), followed by a	TCZ: 8 mg per kilogram of body weight administered	TCZ 8 mg/kg, maximum 800 mg	TCZ 8 mg/kg, maximum 800 mg +standard care per local practice	tocilizumab single intravenous administration on 8mg/Kg	TCZ 8 mg/kg, maximum 800 mg +standard care	TCZ 400 mg +standard care per local practice	tocilizumab+favipiravir Favipiravir: On the 1st day, 1600mg twice a day; from



Author, year, reference number/Study name/Study ID	Rosas et al 2020 [9] COVACTA NCT04320615	Salvarani et al 2020 [10] NCT04346355	Stone et al 2020 [11] NCT04356937	Salama et al 2020 [13] EMPACTA NCT04372186	Hermine et al 2020 [14] NCT04331808	Gordon et al 2021 [15] REMAP-CAP	Veiga et al. 2021 [12] TOCIBRAS	Wang et al 2020 [16] ChiCTR2000029765	Zhao et al. 2020 [17] NCT04310228 ChiCTR2000030894
<b>number of randomized/enrolled patients in subgroups - Mild, Moderate, Severe, Critical COVID-19)</b>	maximum dose 800 mg; up to 1 additional dose may be given if clinical symptoms worsen or show no improvement	second dose after 12 hours	intravenously, not to exceed 800 mg						the 2nd to the 7th day, 600mg twice a day. Oral administration, the maximum number of days taken is not more than 7 days. Tocilizumab: The first IV dose is 4 ~ 8mg/kg and the recommended dose is 400mg. For fever patients, an additional application (the same dose as before) is given if there is still fever within 24 hours after the first dose and the interval between two medications ≥ 12 hours. (mild n=0; common/moderate n=8; severe n=5; critical n=1) <u>tocilizumab</u> (mild n=0; common/moderate n=2; severe n=5; critical n=0)
<b>Comparator (s)</b>	placebo	supportive care following the	placebo	placebo+standard care per local practice	placebo+standard care per local practice	sarilumab single iv	standard care	placebo+standard care per local practice	favipiravir (mild n=0; common/moderate

<b>Author, year, reference number/Study name/Study ID</b>	Rosas et al 2020 [9] COVACTA NCT04320615	Salvarani et al 2020 [10] NCT04346355	Stone et al 2020 [11] NCT04356937	Salama et al 2020 [13] EMPACTA NCT04372186	Hermine et al 2020 [14] NCT04331808	Gordon et al 2021 [15] REMAP-CAP	Veiga et al. 2021 [12] TOCIBRAS	Wang et al 2020 [16] ChiCTR2000029765	Zhao et al. 2020 [17] NCT04310228 ChiCTR2000030894
<b>(standard care or generic drug name and dosage, time frame; number of randomized/enrolled patients in subgroups - Mild, Moderate, Severe, Critical COVID-19)</b>		treatment protocols of each center				administration 400mg			n=2; severe n=3; critical n=0)
<b>Primary Outcome(s)</b>	clinical status on a 7-category ordinal scale at day 28	clinical worsening within 14 days since randomization	intubation (or death, for patients who died before intubation) after administration of tocilizumab or placebo	cumulative proportion of patients requiring mechanical ventilation	>5 on the World Health Organization 10-point Clinical Progression Scale (WHO-CPS) on day 4 and survival without need of ventilation at day 14	an ordinal scale combining in-hospital mortality and days free of organ support to day 21	clinical status measured at 15 days using a seven level ordinal scale	cure rate	cumulative lung lesion remission rate
<b>Patient-relevant secondary outcome(s)</b>	clinical status at day 14 on the 7-category ordinal scale,	evaluation of the efficacy of early vs late administration of tocilizumab in admission	clinical worsening, discontinuation of supplemental oxygen,	time to hospital discharge or ready for discharge; time to a $\geq 2$ category improvement in	clinical status assessed with the WHO-CPS scores at day 7 and day 14, overall survival, time to discharge, time	all-cause mortality; hospital mortality Health-related Quality of life	all-cause mortality hospital mortality improvement of SOFA scale	mortality; ventilator utilization; hospitalization day	improvement of clinical symptoms the changes of blood routine test and IL-6; changes of oxygen therapy

<b>Author, year, reference number/Study name/Study ID</b>	Rosas et al 2020 [9] COVACTA NCT04320615	Salvarani et al 2020 [10] NCT04346355	Stone et al 2020 [11] NCT04356937	Salama et al 2020 [13] EMPACTA NCT04372186	Hermine et al 2020 [14] NCT04331808	Gordon et al 2021 [15] REMAP-CAP	Veiga et al. 2021 [12] TOCIBRAS	Wang et al 2020 [16] ChiCTR2000029765	Zhao et al. 2020 [17] NCT04310228 ChiCTR2000030894
	mortality at day 28, ventilator-free days to day 28, time to improvement from baseline in $\geq 2$ categories on the 7-category ordinal scale, and time to hospital discharge	to ICU with mechanical ventilation, mortality, and tocilizumab toxic effects		clinical status; time to clinical failure; mortality rate	to oxygen supply independency, biological factors such as C-reactive protein level, and adverse events.	assessment, EQ5D-5L and WHODAS 2.0	ventilator free days		
<b>Follow-up (days, months)</b>	60 days	30 days	28 days	60 days	90 days	90 days	29 days		3 months
<b>Sponsor/lead institution</b>	Hoffmann-La Roche	Azienda Unità Sanitaria Locale Reggio Emilia	Massachusetts General Hospital	Genentech, Inc.	Assistance Publique - Hôpitaux de Paris	MJM Bonten, UMC Utrecht	Beneficência Portuguesa de São Paulo	The First Affiliated Hospital of University of science and technology of China (Anhui Provincial Hospital)	Peking University First Hospital

\*Mixed COVID-19, Mild, Moderate, Severe, Critical COVID-19

**Table 4-7 Summary of safety from observational studies (AE and SAE) of tocilizumab**

Author, year	Xu et al 2020 [7]	Luo et al 2020 [17]	Toniati et al 2020 [8]	Somers et al 2020 [21]	Rossi et al 2020 [22]	Petrak et al [23]	Mikulska et al. [24]	Malekzadeh et al. [25]	Perrone et al. [26]	Di Niso et al 2020 [27]	Sciascia et al 2020 [6]	Price et al 2020 [29]
<b>Country</b>	China	China	Italy	USA	France	USA	Italy	Iran	Italy	Italy	Italy	USA
<b>Sponsor/lead institution</b>	Department of Science and Technology of Anhui Province and Health Commission of Anhui Province China National Center for Biotechnology Development 175	Zhongfaxin cheng campus of Tongji Hospital	n.a.	National Institutes of Health Centers for Disease Control and Prevention American Society for Transplantation and Cellular Therapy New Investigator Award	Centre Hospitalier Intercommunal Robert Ballanger Groupe Hospitalier Pitie-Salpetriere	n.a.	n.a.	AryoGen Co., Iran	National Cancer Institute, Naples	n.a.	ASL Città di Torino	The Yale School of Medicine Institutional Review Board (2000027792)
<b>Intervention/Product (drug name)</b>	tocilizumab lopinavir/ritonavir; INF- $\alpha$ ; ribavirin;	tocilizumab/ tocilizumab +methylprednisolone	tocilizumab + standard pharmacological protocol	tocilizumab + standard pharmacological protocol	tocilizumab	tocilizumab + steroid or vasopressors or hydroxychloroquine and azithromycin	tocilizumab and/or methylprednisolone+S OC	tocilizumab	tocilizumab	tocilizumab + SoC	tocilizumab	tocilizumab

<b>Dosage</b>	4-8 mg/kg max 800 mg	n.a.	8 mg/kg max 800 mg	8 mg/kg max 800 mg	400 mg	4 mg/kg max 800 mg	8 mg/kg max 800 mg, methylpred nisolone 1mg/kg for 5 days intravenous ly, then 0.5mg/kg for 5 days	sc 324 mg tocilizuma b <100 kg; ≥100 kg 486 mg + SOC	2 doses of TCZ 8 mg/kg (up to a maximum of 800mg per dose), with an interval of 12 hours	sc 324 mg tocilizuma b + SOC	sc 324 mg, iv 8 mg/kg max 800 mg tocilizu mab	iv 8 mg/kg tocilizuma b
<b>Comparator</b>	n.a.	n.a.	n.a.	standard pharmacolo gical protocol	standard pharmacolo gical protocol	n.a.	standard pharmacolo gical protocol	n.a.	n.a.	n.a.	no	no
<b>Study design</b>	observati onal	observati onal	observati onal	observati onal, controlled study	observati onal	observati onal	observati onal	observati onal	observati onal	observati onal	observati onal	observati onal
<b>Setting</b>	hospital	hospital	hospital	hospital	hospital	hospital	hospital	hospital	hospital	hospital	hospital	hospital
<b>Number of pts</b>	21	15	100	154	246	145	196	126	301	70	63	239
<b>Inclusion criteria</b>	patients with severe and criticalCO VID-19	patients infected with COVID-19	infected with COVID-19; absence of contraindic ation to tocilizumab	patients were admitted to Michigan Medicine from March 9- April 20, 2020 for severe COVID-19 pneumonia ,required invasive mechanical ventilation	patients hospitalized with COVID-19	patients hospitalized with COVID-19	COVID-19 pneumonia	severe or critical COVID- 19	patients hospitalized with COVID-19	COVID- 19 pneumonia	patients infected with COVID- 19	patients infected with COVID- 19

<b>Exclusion criteria</b>	active pulmonary tuberculosis combined with clear bacterial infection and fungal infection	n.a.	contraindication to TCZ, suspected or confirmed bacterial infection, an active diverticulitis or GIT perforation, neutropenia, thrombocytopenia	<16 years intubated for conditions unrelated to COVID-19, enrolled into a RCT for sarilumab	lack of consent of palliative care patients in ICU patients transferred from ICU	missing data	intubated treated with remdesivir pregnancy	hypersensitivity to TCZ, history of infection HBV, HCV, HIV, hepatic disorders, bone marrow suppression, active peptic ulcer, diverticulitis, or any other GI disorders that increase the risk of GI perforation, pregnancy, breastfeeding, any concurrent active infection other than COVID-19, and severe renal impairment	contraindicated TCZ, ALT/AST > 5 times the upper limit of the normality, neutrophils count < 500 /mmc, platelets < 50.000/mmc, bowel diverticulitis or perforation	suspected or confirmed concomitant bacterial or fungal systemic infection, active diverticulitis or GIT perforation, neutropenia, platelets count < 50 x 10 <sup>9</sup> /L, creatinine clearance < 30 ml/min, serum levels of ALT/AST > 5 times the upper limit of the normal range, or pregnancy.	n.a.	n.a.
<b>Age of patients (yrs)</b>	56.8±16.5 (25–88)	73 (62-80)	62 (IQR 57–71])	58±14.9	67.6 ±15.3	58.1	67.9 years (range, 30–100)	median 55 (20–85) years	≤ 60: 122 (40.5%) 61-70: 107 (35.5%) 71 + : 72 (23.9%)	median 60 (IQR 52, 75)	mean	median

<b>Disease severity</b>	severe	moderate/severe	severe	severe	severe	severe	severe	severe or critical	severe	severe	severe	severe/non severe
<b>Follow-up (months)</b>	Hospitalization days (range) 15.1±5.8 (10–31)	1 week after tocilizumab therapy	10-day follow-up	Median follow-up 47 days (28-67).	28-day maximum follow-up	15.3 days length of hospital stay	median follow-up of 53 days (range 4–70, interquartile range 33–57)	median 8 days (5–12)	up to 30 days	35 days	14 days	≥21 days
<b>Loss to follow-up, n (%)</b>	0	0	0	0	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
<b>RoB</b>	high	high	high	high	high	high	high	high	high	high	high	high
<b>Safety – Outcomes*</b>												
<b>Overall AEs, n (%)</b>	n.a.	n.a.	100%	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
<b>Serious AE (SAE), n (%)</b>	0%	n.a.	3%	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
<b>Most frequent AEs n (%)</b>	n.a.	n.a.	100%	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
<b>Most frequent SAEs, n (%)</b>	n.a.	n.a.	3%	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
<b>AEs of special interest, n (%)</b>	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
<b>Death as SAE, n (%)</b>	n.a.	n.a.	2%	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
<b>Withdrawals due AEs, n (%)</b>	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.

\* by arms, if available, (Robins-I):

<https://training.cochrane.org/handbook/current/chapter-25>

Abbreviations: CI=Confidence interval; RR=Risk ratio

**Table 4-8 Ongoing trials of single agent tocilizumab**

<b>Trial Identifier/registry ID(s)/contact</b>	ChiCTR2000029765	NCT04317092 TOCIDVID-19	NCT04306705 TACOS	NCT04315480	NCT04372186 EMPACTA	NCT02735707 REMAP-CAP	NCT04381936 RECOVERY
<b>Study design, study phase</b>	Phase 4 A multicenter, randomized controlled trial for the efficacy and safety of tocilizumab in the treatment of new coronavirus pneumonia (COVID-19). RCT parallel	Phase 2 Multicenter single-arm, open-label, phase 2 study on the efficacy and tolerability of tocilizumab in the treatment of patients with COVID-19 pneumonia non randomized	A Retrospective Study of Evaluating Safety and Efficacy of Tocilizumab Compared to Continuous Renal Replacement Therapy in Controlling CRS Triggered by COVID-19 retrospective	Phase 2 Tocilizumab (RoActemra) as Early Treatment of Patients Affected by SARS-CoV2 (COVID-19) Infection With Severe Multifocal Interstitial Pneumonia non randomized, single arm	Phase III A Randomized, Double-Blind, Placebo-Controlled, Multicenter Study to Evaluate the Efficacy and Safety of Tocilizumab in Hospitalized Patients With COVID-19 Pneumonia RCT parallel	Phase IV randomized, open label; international, multifactorial, adaptive platform trial	Phase 2 and 3 randomized, open label; factorial assignment
<b>Recruitment status</b>	Recruiting	Active, not recruiting	Recruiting	Active, not recruiting	Active, not recruiting	Recruiting	Recruiting
<b>Number of Patients, Disease severity*</b>	198 severe	301 n.a.	target sample size: 120 n.a.	38 severe	379.n.a.	353 (toc vs, sarilumab) patients critically ill COVID-19.	2022 severe participants with progressive COVID-19
<b>Setting (hospital, ambulatory,..)</b>	Hospital	Hospital	Hospital	Hospital	Hospital	hospital	hospital
<b>Intervention drug name and dosage</b>	tocilizumab, n.a	tocilizumab 2 doses of TCZ 8 mg/kg (up to a maximum of 800mg per dose), with an interval of 12 hours	Tocilizumab or CRRT (continuous renal replacement therapy) or SoC	tocilizumab single intravenous administration 8mg/Kg	tocilizumab 8 mg/kg IV (max 800 mg) + SOC	tocilizumab single intravenous administration 8mg/Kg	tocilizumab 8 mg/kg IV (max 800 mg).
<b>Comparator (drug name and dosage)</b>	conventional therapy	n.a.	n.a.	n.a.	placebo+ SOC	sarilumab single iv administration 400mg	no additional treatment
<b>Primary Outcomes</b>	cure rate mortality;ventilator	Lethality rate two weeks / one	Proportion of Participants With Normalization of	arrest in deterioration of pulmonary	Cumulative Proportion of Participants	respiratory and cardiovascular	all-cause mortality



<b>Trial Identifier/registry ID(s)/contact</b>	ChiCTR2000029765	NCT04317092 TOCIDVID-19	NCT04306705 TACOS	NCT04315480	NCT04372186 EMPACTA	NCT02735707 REMAP-CAP	NCT04381936 RECOVERY
	utilization;hospitalization day	month after registration	Fever and Oxygen Saturation Through Day 14	function; improving in pulmonary function; need of oro-tracheal intubation; death	Requiring Mechanical Ventilation by Day 28	organ support-free days up to day 21	
<b>Sponsor/ lead institution, country (also country of recruitment if different)</b>	The First Affiliated Hospital of University of science and technology of China (Anhui Provincial Hospital) China	National Cancer Institute, Naples Italy	Tongji Hospital China	Università Politecnica delle Marche Italy	Genentech, Inc. USA, Brazil, Kenya, Mexico, Peru, South Africa	MJM Bonten, UMC Utrecht	University of Oxford

**Table 4-9 Ongoing trials of combination therapies tocilizumab**

<b>Trial Identifier</b>	NCT04332094	NCT04424056	NCT04409262 REMDACTA	NCT04310228	NCT04600141	NCT04330638 COV-AID	NCT04678739	NCT04779047
<b>Study design, study phase</b>	Phase 2, Pilot, Randomized, Multicenter, Open-label RCT parallel, open label	Phase 3 open label randomized therapeutic trial RCT parallel, open label	Phase III, Randomized, Double-Blind, Multicenter Study RCT parallel, double blind	Not Applicable Phase, Multicenter, RCT parallel, open label	Phase III, Parallel Assignment RCT parallel, open label	Phase III; randomized, open label	Phase III; randomized, open label	Phase 4 randomized, open label
<b>Recruitment status</b>	Recruiting	Not yet recruiting	Active, not recruiting	Recruiting	Not yet recruiting	Active, not recruiting	Recruiting	Recruiting
<b>Number of Patients, Disease severity</b>	276 patients Severity 3-4 according to the WHO 7-point ordinal scale	216 patients COVID19 infection pneumonia at stage 2b or advanced stage 3	450 patients severe COVID-19 pneumonia	150 patients n.a.	severe COVID-19 infection	342, critical ill	150 patients severe COVID-19.	150 patients n.a.
<b>Setting (hospital, ambulatory,..)</b>	hospital	hospital	hospital	hospital	hospital	hospital	hospital	hospital
<b>Intervention drug name and dosage</b>	tocilizumab, hydroxychloroquine, azithromycin	anakinra +/- ruxolitinib tocilizumab +/- ruxolitinib	remdesivir + tocilizumab	favipiravir + tocilizumab	Tocilizumab iv 8mg/kg/dose + Therapeutic dosage heparin	tocilizumab (iv 8mg/kg/dose) tocilizumab+anakinra (100 mg for 28 days) anakinra anakinra+siltuximab (iv 11 mg/kg) siltuximab	A loading dose of Remdesivir I/V 5mg/kg (less than 40kg) or 200mg (>40kg) on day 1, then 2.5mg/kg (less than 40kg) or 100mg (>40kg) daily following randomization. Tocilizumab I/V 8mg/Kg up to 800mg highest 12 hours apart	remdesivir iv 200 mg at day 1 then 100 mg once daily for 5 days and tocilizumab 800 mg once
<b>Comparator (drug name and dosage)</b>	hydroxychloroquine, azithromycin	standard of care	remdesevir	favipiravir, tocilizumab	Heparin - Therapeutic dosage (Group 1) and Heparin -	usual care	Treatment as given without Remdesivir and Tocilizumab	hydroxychloroquine 400 mg twice daily at day 1 then 200 mg twice daily for 5 days and

<b>Trial Identifier</b>	NCT04332094	NCT04424056	NCT04409262 REMDACTA	NCT04310228	NCT04600141	NCT04330638 COV-AID	NCT04678739	NCT04779047
					Prophylactic dosage (Group 2)			tocilizumab 800 mg once
<b>Primary Outcomes</b>	in-hospital mortality	ventilation free days at D28	Clinical Status as Assessed by the Investigator Using a 7-Category Ordinal Scale of Clinical Status on Day 2	Clinical cure rate	Proportion of patients with clinical improvement in 30 days	Time to clinical improvement	Time to clinical improvement Duration of ICU Stay; Mortality Rate; Time to Recovery, Hospital stay	percentage of clinical cure
<b>Sponsor/ lead institution, country (also country of recruitment if different)</b>	Fundació Institut de Recerca de l'Hospital de la Santa Creu i Sant Pau Spain	Assistance Publique Hopitaux De Marseille France	Hoffmann-La Roche Brazil, Russian Federation, United States	Peking University First Hospital China	University of Sao Paulo Brazil	University Hospital, Ghent Belgium	M Abdur Rahim Medical College and Hospital	October 6 University

## 5 REFERENCES

- [1] Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.0 (updated July 2019). Cochrane, 2019. Available from <http://www.training.cochrane.org/handbook>.
- [2] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986; **7**: 177-188.
- [3] Balshem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, Vist GE, Falck-Ytter Y, Meerpohl J, Norris S, Guyatt GH. GRADE guidelines: 3. Rating the quality of evidence. *Journal of Clinical Epidemiology* 2011; **64**: 401-406.
- [4.] Yoshikawa T., Hill T., Li K., Peters C.J., Tseng C.T. Severe acute respiratory syndrome (SARS) coronavirus-induced lung epithelial cytokines exacerbate SARS pathogenesis by modulating intrinsic functions of monocyte-derived macrophages and dendritic cells. *J Virol*. 2009;**83**(7):3039-3048. doi:10.1128/JVI.01792-08.
- [5.] European Medicines Agency. EMA. RoActemra (tocilizumab) 2020 [Available from: <https://www.ema.europa.eu/en/medicines/human/EPAR/roactemra>
- [6.] Sciascia S., Aprà F., Baffa A., al. e. Pilot prospective open, single-arm multicentre study on off-label use of tocilizumab in patients with severe COVID-19. *Clin Exp Rheumatol*. 2020;**38**(3):529-32
- [7.] Xu X., Han M., Li T., Sun W., Dongsheng W., Fu B., et al. Effective Treatment of Severe COVID-19 Patients with Tocilizumab. *Proc Natl Acad Sci U S A*. 2020;**117**(20):10970-10975. doi:10.1073/pnas.2005615117
- [8.] Toniati P, Piva S, Cattalini M, Garrafa E, Regola F, Castelli F, Franceschini F, Airò P, Bazzani C, Beindorf EA, Berlendis M, Bezzi M, Bossini N, Castellano M, Cattaneo S, Cavazzana I, Contessi GB, Crippa M, Delbarba A, De Peri E, Faletti A, Filippini M, Filippini M, Frassi M, Gaggiotti M, Gorla R, Lanspa M, Lorenzotti S, Marino R, Maroldi R, Metra M, Matteelli A, Modena D, Moioli G, Montani G, Muiesan ML, Odolini S, Peli E, Pesenti S, Pezzoli MC, Pirola I, Pozzi A, Proto A, Rasulo FA, Renisi G, Ricci C, Rizzoni D, Romanelli G, Rossi M, Salvetti M, Scolari F, Signorini L, Taglietti M, Tomasoni G, Tomasoni LR, Turla F, Valsecchi A, Zani D, Zuccalà F, Zunica F, Focà E, Andreoli L, Latronico N. Tocilizumab for the treatment of severe COVID-19 pneumonia with hyperinflammatory syndrome and acute respiratory failure: A single center study of 100 patients in Brescia, Italy. *Autoimmun Rev*. 2020 Jul;**19**(7):102568. doi: 10.1016/j.autrev.2020.102568. Epub 2020 May 3. PMID: 32376398; PMCID: PMC7252115.
- [9.] Rosas I, Bräu N, Waters M, Go RC, Hunter BD, Bhagani S, et al. Tocilizumab in Hospitalized Patients With COVID-19 Pneumonia. *medRxiv*. 2020:2020.08.27.20183442.
- [10.] Salvarani C, Dolci G, Massari M, Merlo DF, Cavuto S, Savoldi L, et al; RCT-TCZ-COVID-19 Study Group. Effect of Tocilizumab vs Standard Care on Clinical Worsening in Patients Hospitalized With COVID-19 Pneumonia: A Randomized Clinical Trial. *JAMA Intern Med*. 2020 Oct 20:e206615. doi: 10.1001/jamainternmed.2020.6615. Epub ahead of print. PMID: 33080005; PMCID: PMC7577199.
- [11.] Stone JH, Frigault MJ, Serling-Boyd NJ, Fernandes AD, Harvey L, Foulkes AS, et al; BACC Bay Tocilizumab Trial Investigators. Efficacy of Tocilizumab in Patients Hospitalized with Covid-19. *N Engl J Med*. 2020 Oct 21. doi: 10.1056/NEJMoa2028836. Epub ahead of print. PMID: 33085857.

- [12.] Veiga VC, Prats JAGG, Farias DLC, Rosa RG, Dourado LK, Zampieri FG, et al. Effect of tocilizumab on clinical outcomes at 15 days in patients with severe or critical coronavirus disease 2019: randomised controlled trial. *BMJ*. 2021;372:n84.
- [13.] Salama C, Han J, Yau L, Reiss WG, Kramer B, Neidhart JD, et al. Tocilizumab in nonventilated patients hospitalized with Covid-19 pneumonia. *medRxiv*. 2020:2020.10.21.20210203.
- [14.] Hermine O, Mariette X, Tharaux PL, Resche-Rigon M, Porcher R, Ravaud P; CORIMUNO-19 Collaborative Group. Effect of Tocilizumab vs Usual Care in Adults Hospitalized With COVID-19 and Moderate or Severe Pneumonia: A Randomized Clinical Trial. *JAMA Intern Med*. 2020 Oct 20:e206820. doi: 10.1001/jamainternmed.2020.6820. Epub ahead of print. PMID: 33080017; PMCID: PMC7577198
- [15.] Gordon AC, Mouncey PR, Al-Beidh F, Rowan KM, Nichol AD, Arabi YM, et al. Interleukin-6 Receptor Antagonists in Critically Ill Patients with Covid-19 – Preliminary report. *medRxiv*. 2021:2021.01.07.21249390.
- [16.] Wang D, Fu B, Peng Z, Yang D, Han M, Li M, et al. Tocilizumab Ameliorates the Hypoxia in COVID-19 Moderate Patients with Bilateral Pulmonary Lesions: A Randomized, Controlled, Open-Label, Multicenter Trial.. *The Lancet*. 2020 . Available at SSRN: <https://ssrn.com/abstract=3667681> or <http://dx.doi.org/10.2139/ssrn.3667681>
- [17.] Zhao H, Zhu Q, Zhang C, Li J, Wei M, Qin Y, et al. Tocilizumab combined with favipiravir in the treatment of COVID-19: A multicenter trial in a small sample size. *Biomedicine & Pharmacotherapy*. 2021;133:110825. doi.org/10.1016/j.biopha.2020.110825.
- [18.] Cruciani F, De Crescenzo F., Vecchi S., Saulle R., Mitrova Z., Amato L., et al. Efficacy of Early Administration of Tocilizumab in COVID-19 Patients. 2020 [Available from: <https://www.aifa.gov.it/web/guest/-/covid-19-studio-randomizzato-italiano-nessun-beneficio-dal-tocilizumab>
- [19.] Luo P, Liu Y, Qiu L, Liu X, Liu D, Li J. Tocilizumab treatment in COVID-19: A single center experience. *J Med Virol*. 2020 Jul;92(7):814-818. doi: 10.1002/jmv.25801. Epub 2020 Apr 15. PMID: 32253759; PMCID: PMC7262125.
- [20.] Xu X., Han M., Li T., Sun W., Dongsheng W., Fu B., et al. Effective Treatment of Severe COVID-19 Patients with Tocilizumab. *Proc Natl Acad Sci U S A*. 2020;117(20):10970-10975. doi:10.1073/pnas.2005615117 Xu X., Han M., Li T., Sun W., Dongsheng W., Fu B., et al. Effective Treatment of Severe COVID-19 Patients with Tocilizumab. *Proc Natl Acad Sci U S A*. 2020;117(20):10970-10975. doi:10.1073/pnas.2005615117
- [21.] Somers EC, Eschenauer GA, Troost JP, Golob JL, Gandhi TN, Wang L, Zhou N, Petty LA, Baang JH, Dillman NO, Frame D, Gregg KS, Kaul DR, Nagel J, Patel TS, Zhou S, Luring AS, Hanauer DA, Martin E, Sharma P, Fung CM, Pogue JM. Tocilizumab for treatment of mechanically ventilated patients with COVID-19. *Clin Infect Dis*. 2020 Jul 11:ciaa954. doi: 10.1093/cid/ciaa954. Epub ahead of print. PMID: 32651997; PMCID: PMC7454462.
- [22.] Rossi B, Nguyen LS, Zimmermann P, Boucenna F, Dubret L, Baucher L, Guillot H, Bouldouyre MA, Allenbach Y, Salem JE, Barsoum P, Oufella A, Gros H. Effect of Tocilizumab in Hospitalized Patients with Severe COVID-19 Pneumonia: A Case-Control Cohort Study. *Pharmaceuticals (Basel)*. 2020 Oct 17;13(10):317. doi: 10.3390/ph13100317. PMID: 33080877; PMCID: PMC7603074.

- [23.] Petrak RM, Skorodin NC, Van Hise NW, Fliegelman RM, Pinsky J, Didwania V, Anderson M, Diaz M, Shah K, Chundi VV, Hines DW, Harting BP, Sidwha K, Yu B, Brune P, Owaisi A, Beezhold D, Kent J, Vais D, Han A, Gowda N, Sahgal N, Silverman J, Stake J, Nepomuceno J, Heddurshetti R. Tocilizumab as a Therapeutic Agent for Critically Ill Patients Infected with SARS-CoV-2. *Clin Transl Sci*. 2020 Sep 12. doi: 10.1111/cts.12894. Epub ahead of print. PMID: 32918792.
- [24.] Mikulska M, Nicolini LA, Signori A, Di Biagio A, Sepulcri C, Russo C, Dettori S, Berruti M, Sormani MP, Giacobbe DR, Vena A, De Maria A, Dentone C, Taramasso L, Mirabella M, Magnasco L, Mora S, Delfino E, Toscanini F, Balletto E, Alessandrini AI, Baldi F, Briano F, Camera M, Dodi F, Ferrazin A, Labate L, Mazzarello G, Pincino R, Portunato F, Tutino S, Barisione E, Bruzzzone B, Orsi A, Schenone E, Rossetti N, Sasso E, Da Rin G, Pelosi P, Beltramini S, Giacomini M, Icardi G, Gratarola A, Bassetti M. Tocilizumab and steroid treatment in patients with COVID-19 pneumonia. *PLoS One*. 2020 Aug 20;15(8):e0237831. doi: 10.1371/journal.pone.0237831. PMID: 32817707; PMCID: PMC7440633.
- [25.] Malekzadeh R, Abedini A, Mohsenpour B et al. Subcutaneous tocilizumab in adults with severe and critical COVID-19: A prospective open-label uncontrolled multicenter trial. *Int Immunopharmacol*. 2020 Oct 13;89(Pt B):107102. doi: 10.1016/j.intimp.2020.107102. Epub ahead of print. PMID: 33075713; PMCID: PMC7553010.
- [26.] Perrone F, Piccirillo MC, Ascierio PA, et al.; TOCIVID-19 investigators, Italy. Tocilizumab for patients with COVID-19 pneumonia. The single-arm TOCIVID-19 prospective trial. *J Transl Med*. 2020 Oct 21;18(1):405. doi: 10.1186/s12967-020-02573-9. PMID: 33087150; PMCID: PMC7576974.
- [27.] Di Nisio M, Potere N, Candeloro M, Spacone A, Pieramati L, Ferrandu G, Rizzo G, La Vella M, Di Carlo S, Cibelli D, Parruti G, Levi M, Porreca E. Interleukin-6 receptor blockade with subcutaneous tocilizumab improves coagulation activity in patients with COVID-19. *Eur J Intern Med*. 2020 Nov 3:S0953-6205(20)30406-4. doi: 10.1016/j.ejim.2020.10.020. Epub ahead of print. PMID: 33162299; PMCID: PMC7608031.
- [28.] F Hoffman-La Roche Ltd. Roche's phase III EMPACTA study showed Actemra/RoActemra reduced the likelihood of needing mechanical ventilation in hospitalised patients with COVID-19 associated pneumonia. Published September 18, 2020. Accessed September 19, 2020. <https://www.roche.com/investors/updates/inv-update-2020-09-18.htm>
- [27.] Price CC, Altice FL, Shyr Y, Koff A, Pischel L, Goshua G, Azar MM, Mcmanus D, Chen SC, Gleeson SE, Britto CJ, Azmy V, Kaman K, Gaston DC, Davis M, Burrello T, Harris Z, Villanueva MS, Aoun-Barakat L, Kang I, Seropian S, Chupp G, Bucala R, Kaminski N, Lee AI, LoRusso PM, Topal JE, Dela Cruz C, Malinis M. Tocilizumab Treatment for Cytokine Release Syndrome in Hospitalized Patients With Coronavirus Disease 2019: Survival and Clinical Outcomes. *Chest*. 2020 Oct;158(4):1397-1408. doi: 10.1016/j.chest.2020.06.006. Epub 2020 Jun 15. PMID: 32553536; PMCID: PMC7831876.
- [30.] Live map of COVID-19 evidence [web page]. Oslo: Norwegian Institute of Public Health. [updated 21. January 2021; cited 09. February 2021]. Available from: <https://www.fhi.no/en/qk/systematic-reviews-hta/map/>
- [31.] COVID-19: a living systematic map of the evidence [web page]. London: EPPI Centre, University College London. [updated 04. February 2021; cited 09. February 2021]. Available from: <http://eppi.ioe.ac.uk/cms/Projects/DepartmentofHealthandSocialCare/Publishedreviews/COVID-19Livingssystematicmapofthevidence/tabid/3765/Default.aspx>

## 6 APPENDIX

### 6.1 Search strategy to identify randomised controlled trials

DEPLazio, the Department of Epidemiology of the Regional Health Service Lazio in Rome, Italy is responsible for setting up the search strategy to identify randomised controlled trials (RCTs). DEPLazio performed a search in Medline, PubMed, and Embase, which has been updated weekly from March 2020 (Appendix Table 6-1). DEPLazio searched medRxiv.org (<https://www.medrxiv.org/>), bioRxiv.org (<https://www.biorxiv.org/>), and arXiv.org (<https://www.arxiv.org/>) for preprints of preliminary reports of randomised trials. The Cochrane Covid-19 Study Register (<https://covid-19.cochrane.org/>), ClinicalTrials.gov ([www.clinicaltrials.gov](http://www.clinicaltrials.gov)) and World Health Organization (WHO) International Clinical Trials Registry Platform (ICTRP) ([www.who.int/ictcp/en/](http://www.who.int/ictcp/en/)) were search in addition. Other sources included journal alerts, contact with researchers, websites such as Imperial College, London School of Hygiene and Tropical Medicine, and Eurosurveillance. We applied no restriction on language of publication.

We included randomised controlled trials (RCTs) comparing any pharmacological intervention against another pharmacological intervention or placebo or standard care (SC), for the treatment of individuals with Covid-19. We excluded studies comparing two dosages of the same pharmacological agent. We did not exclude studies on individuals with a comorbid disorder.

Four authors independently screened the references retrieved by the search, selected the studies, and extracted the data, using a predefined data-extraction sheet. The same reviewers discussed any uncertainty regarding study eligibility and data extraction until consensus was reached; conflicts of opinion were resolved with other members of the review team. Two authors independently assessed the risk of bias of the included studies with the Cochrane tool. Three authors used the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach, to evaluate the strength of evidence.

The methods described above are part of a living review of pharmacological agents for the treatment of Covid-19 conducted by the Department of Epidemiology of the Regional Health Service Lazio, Italy, to inform national regulatory agencies and clinicians, available at <https://www.deplazio.net/farmacicovid>. The review is registered on Prospero (CRD42020176914).

**Table 6-1 Search strategy to identify randomised controlled studies**

Database	URL	Search line / Search terms	Date of search
Pubmed	pubmed.ncbi.nlm.nih.gov	<p>1. (((((((("Coronavirus"[Mesh]) OR (coronavirus*[Title/Abstract] OR coronavirus*[Title/Abstract] OR coronavirus*[Title/Abstract] OR coronavirus*[Title/Abstract] OR Wuhan*[Title/Abstract] OR Hubei*[Title/Abstract] OR Huanan[Title/Abstract] OR "2019-nCoV"[Title/Abstract] OR 2019nCoV[Title/Abstract] OR nCoV2019[Title/Abstract] OR "nCoV-2019"[Title/Abstract] OR "COVID-19"[Title/Abstract] OR COVID19[Title/Abstract] OR "CORVID-19"[Title/Abstract] OR CORVID19[Title/Abstract] OR "WN-CoV"[Title/Abstract] OR WNCov[Title/Abstract] OR "HCoV-19"[Title/Abstract] OR HCoV19[Title/Abstract] OR CoV[Title/Abstract] OR "2019 novel"[Title/Abstract] OR Ncov[Title/Abstract] OR "n-cov"[Title/Abstract] OR "SARS-CoV-2"[Title/Abstract] OR "SARSCoV-2"[Title/Abstract] OR "SARSCoV2"[Title/Abstract] OR "SARS-CoV2"[Title/Abstract] OR SARSCov19[Title/Abstract] OR "SARS-Cov19"[Title/Abstract] OR "SARSCov-19"[Title/Abstract] OR Ncovor[Title/Abstract] OR Ncorona*[Title/Abstract] OR Ncorono*[Title/Abstract] OR NcovWuhan*[Title/Abstract] OR NcovHubei*[Title/Abstract] OR NcovChina*[Title/Abstract] OR NcovChinese*[Title/Abstract])) OR (((respiratory*[Title/Abstract] AND (symptom*[Title/Abstract] OR disease*[Title/Abstract] OR illness*[Title/Abstract] OR condition*[Title/Abstract] OR "seafood market"[Title/Abstract] OR "food market"[Title/Abstract] AND (Wuhan*[Title/Abstract] OR Hubei*[Title/Abstract] OR China*[Title/Abstract] OR Chinese*[Title/Abstract] OR Huanan*[Title/Abstract])) OR ("severe acute respiratory syndrome")) OR ((corona*[Title/Abstract] OR corono*[Title/Abstract] AND (virus*[Title/Abstract] OR viral*[Title/Abstract] OR virinae*[Title/Abstract])) AND (((((((randomized controlled trial [pt]) OR (controlled clinical trial [pt]) OR (randomized [tiab]) OR (placebo [tiab]) OR (clinical trials as topic [mesh: noexp]) OR (randomly [tiab]) OR (trial [ti])))) NOT (animals [mh] NOT humans [mh]) AND (2019/10/01:2020[dp])</p>	28/02/2020



Database	URL	Search line / Search terms	Date of search
Ovid MEDLINE(R) ALL)	ovidsp.dc2.ovid.com	<ol style="list-style-type: none"> <li>1. exp coronavirus/</li> <li>2. ((corona* or corono*) adj1 (virus* or viral* or virinae*)).ti,ab,kw.</li> <li>3. (coronavirus* or coronovirus* or coronavirinae* or Coronavirus* or Coronovirus* or Wuhan* or Hubei* or Huanan or "2019-nCoV" or 2019nCoV or nCoV2019 or "nCoV-2019" or "COVID-19" or COVID19 or "CORVID-19" or CORVID19 or "WN-CoV" or WNCov or "HCoV-19" or HCoV19 or CoV or "2019 novel*" or Ncov or "n-cov" or "SARS-CoV-2" or "SARSCoV-2" or "SARSCoV2" or "SARS-CoV2" or SARSCov19 or "SARS-Cov19" or "SARSCov-19" or "SARS-Cov-19" or Ncovor or Ncorona* or Ncorono* or NcovWuhan* or NcovHubei* or NcovChina* or NcovChinese*).ti,ab,kw.</li> <li>4. (((respiratory* adj2 (symptom* or disease* or illness* or condition*)) or "seafood market*" or "food market*") adj10 (Wuhan* or Hubei* or China* or Chinese* or Huanan*)).ti,ab,kw.</li> <li>5. ((outbreak* or wildlife* or pandemic* or epidemic*) adj1 (China* or Chinese* or Huanan*)).ti,ab,kw.</li> <li>6. "severe acute respiratory syndrome*".ti,ab,kw.</li> <li>7. or/1-6</li> <li>8. randomized controlled trial.pt.</li> <li>9. controlled clinical trial.pt.</li> <li>10. random*.ab.</li> <li>11. placebo.ab.</li> <li>12. clinical trials as topic.sh.</li> <li>13. random allocation.sh.</li> <li>14. trial.ti.</li> <li>15. or/8-14</li> <li>16. exp animals/ not humans.sh.</li> <li>17. 15 not 16</li> <li>18. 7 and 17</li> <li>19. limit 18 to yr="2019 -Current"</li> </ol>	28/02/2020
OVID EMBASE	ovidsp.dc2.ovid.com	<ol style="list-style-type: none"> <li>1. exp Coronavirinae/ or exp Coronavirus/ exp Coronavirus infection/</li> <li>3. (((("Corona virinae" or "corona virus" or Coronavirinae or coronavirus or COVID or nCoV) adj4 ("19" or "2019" or novel or new)) or (("Corona virinae" or "corona virus" or Coronavirinae or coronavirus or COVID or nCoV) and (wuhan or china or chinese)) or "Corona virinae19" or "Corona virinae2019" or "corona virus19" or "corona virus2019" or Coronavirinae19 or Coronavirinae2019 or coronavirus19 or coronavirus2019 or COVID19 or COVID2019 or nCOV19 or nCOV2019 or "SARS Corona virus 2" or "SARS Coronavirus 2" or "SARS-COV-2" or "Severe Acute Respiratory Syndrome Corona virus 2" or "Severe Acute Respiratory Syndrome Coronavirus 2").ti,ab,kw.</li> <li>4. or/1-3</li> <li>5. Clinical-Trial/ or Randomized-Controlled-Trial/ or Randomization/ or Single-Blind-Procedure/ or Double-Blind-Procedure/ or Crossover-Procedure/ or Prospective-Study/ or Placebo/</li> <li>6. (((clinical or control or controlled) adj (study or trial)) or ((single or double or triple) adj (blind\$3 or mask\$3)) or (random\$ adj (assign\$ or allocat\$ or group or grouped or patients or study or trial or distribut\$)) or (crossover adj (design or study or trial)) or placebo or placebos).ti,ab.</li> <li>7. 5 or 6</li> <li>8. 4 and 7</li> <li>9. limit 8 to yr="2019 -Current"</li> </ol>	28/02/2020

## 6.2 Search strategy to identify observational studies

As of October 2020, NIPHNO is responsible for setting up the search strategy to identify observational studies.

From September to December 2020, we received records that EPPI Centre has screened after searching weekly in Medline and Embase (until beginning of November 2020), from November onwards Microsoft Academic Graph (MAG). We supplemented these studies with a weekly search in Scopus (Elsevier). Detailed descriptions of the EPPI and NIPHNO searches are given at their websites [30];[31]. The retrieved hits were imported to a reference management tool, Endnote (Clarivate Analytics), for deduplication. We then searched the EndNote database using the generic names and synonyms for the included COVID-19 drugs.

From January onwards, an information specialist at NIPHNO has conducted searches in Medline (Ovid), Embase (Ovid) and Scopus (Elsevier) using the search strategy described in table 6.2. To screen the references, two reviewers use a binary machine learning (ML) classifier. References that scored above the identified threshold of 30% certainty to be relevant were retained for screening; while those scoring below this threshold score were set aside.

Prior to using the binary ML classifier score to discard low scoring records, we screened 1028 references manually to train the classifier. The classifier is continuously being updated a long with new references being screened. References that have been set aside, can potentially be picked up in a later stage by a new classifier version. For drugs that have less than 5 publications included in the training batch, we combine the classifier with manual text word searches.

**Table 6-2 Search strategy to identify observational studies**

Database	URL	Search terms / Search modality	Date of search
Embase 1974 to 2021		Lines 1 and 2 are copies of Ovid's Expert searches for covid-19 in MEDLINE and Embase	From 4/2/2021 until 26/2/2021
Ovid MEDLINE(R) ALL 1946 to 2021		<p>1 (((((pneumonia or covid* or coronavirus* or corona virus* or ncov* or 2019-ncov or sars*).mp. or exp pneumonia/) and Wuhan.mp.) or (2019-ncov or ncov19 or ncov-19 or 2019-novel CoV or sars-cov2 or sars-cov-2 or sarscov2 or sarscov-2 or Sars-coronavirus2 or Sars-coronavirus-2 or SARS-like coronavirus* or coronavirus-19 or covid19 or covid-19 or covid 2019 or ((novel or new or nouveau) adj2 (CoV or nCoV or covid or coronavirus* or corona virus or pandemi*2)) or ((covid or covid19 or covid-19) and pandemic*2) or (coronavirus* and pneumonia)).mp. or COVID-19.rx,px,ox,sh. or severe acute respiratory syndrome coronavirus 2.os.) use medall [COVID-19 in MEDLINE]</p> <p>2 (((((pneumonia or covid* or coronavirus* or corona virus* or ncov* or 2019-ncov or sars*).mp. or exp pneumonia/) and Wuhan.mp.) or (coronavirus disease 2019 or 2019-ncov or ncov19 or ncov-19 or 2019-novel CoV or severe acute respiratory syndrome coronavirus 2 or sars-cov2 or sars-cov-2 or sarscov2 or sarscov-2 or Sars-coronavirus2 or Sars-coronavirus-2 or SARS-like coronavirus* or coronavirus-19 or covid19 or covid-19 or covid 2019 or ((novel or new or nouveau) adj2 (CoV or nCoV or covid or coronavirus* or corona virus or pandemi*2)) or ((covid or covid19 or covid-19) and pandemic*2) or (coronavirus* and pneumonia)).mp. or (coronavirus disease 2019 or severe acute respiratory syndrome coronavirus 2).sh,dj.) use oomezd [COVID-19 in Embase]</p> <p>3 (COVID-19 serotherapy/ or Immunization, Passive/ or tocilizumab/ or camostat/ or nafamostat/ or AP301 peptide/ or Interleukin 1 Receptor Antagonist Protein/ or alunacedase alfa/ or darunavir/ or favipiravir/ or sarilumab/ or exp Interferons/ or</p>	

		<p>gimsilumab/ or canakinumab/ or REGN-COV-2/ or bamlanivimab/ or baricitinib/ or molnupiravir/ or Aspirin/ or mavrilimumab/ or exp Vitamin D/dt or Vitamin D Deficiency/dt or Ivermectin/) use medall [MeSH-terms for drugs in MEDLINE]</p> <p>4 (Hyperimmune globulin/dt or tocilizumab/ or camostat/ or camostat mesilate/ or nafamstat/ or solnatide/ or anakinra/ or darunavir/ or favipiravir/ or sarilumab/ or exp Interferon/ or gimsilumab/ or canakinumab/ or baricitinib/ or acetylsalicylic acid/ or mavrilimumab/ or exp Vitamin D/dt or Vitamin D Deficiency/dt or ivermectin/) use oomezd [Emtree-terms for drugs in Embase]</p> <p>5 ((convalescent adj (plasma or sera or serum)) or serotherap* or ((atoxin or hyperimmunoglobulin or hyperimmune globulin or hyperimmune gammaglobulin) adj therap*) or passive immuni?ation or (tocilizumab or atlizumab or (MRA adj monoclonal antibod*) or MSB-11456 or MSB11456 or R-1569 or R1569 or RO-4788533 or RO4788533 or Actemra or Roactemra) or (camostat* or FOY-305 or FOY305 or FOY S 980) or (nafamostat or nafamstat or FUT-175 or FUT175) or (solnatide or AP301 or AP-301 or (TIP adj peptide)) or (anakinra or ((interleukin 1 or IL1 or IL-1) adj2 (antagonist or block* or inhibitor*)) or IL-1Ra or Kineret) or (alunacedase or APN01 or APN-01 or rhACE2 or recombinant human angiotensin converting enzyme 2 or GSK-2586881 or GSK2586881) or (darunavir or prezista or TMC-114 or TMC114 or UIC-94017 or UIC94017) or (favipi?avir or T-705 or T705 or Avigan or Olumiant) or (sarilumab or REGN-88 or REGN88 or SAR-153191 or SAR153191 or Kevzara) or (interferon* or (IFN adj1 (alpha* or beta* or gamma*)) or novaferon or CL-884 or CL884) or (gimsilumab or KIN-1901 or KIN1901 or morab-022 or morab022) or (canakinumab or ACZ-885 or ACZ885 or immunoglobulin G1 or Ilaris) or (REGN-CoV-2 or REGN-CoV2 or antibody cocktail or ((casirivimab or REGN-10933 or REGN10933) and (imdevimab or REGN-10987 or REGN10987))) or (bamlanivimab or LY-CoV555 or LYCoV555 or LY-3819253 or LY3819253) or (baricitinib or LY-3009104 or LY3009104 or INCB-028050 or INCB028050 or INCB-28050 or INCB28050 or Olumiant) or (molnupiravir or MK-4482 or MK4482 or EIDD-2801 or EIDD2801) or (aspirin or acetylsalicylic acid) or (mavrilimumab or immunoglobulin G4 or CAM-3001 or CAM3001) or ((vitamin? D? or D?-vitamin?) adj4 (high-dose* or highdose* or supplement*)) or (ivermect* or MK-933 OR MK933)).mp, bt, ot, du, dy, tn, nm. [other terms (title, abstract, author keywords and more) in MEDLINE and Embase]</p> <p>6 (202009* or 202010* or 202011 or 202012* or 202101* or 202102*).dt. use medall [time limits in MEDLINE]</p> <p>7 (202009* or 202010* or 202011 or 202012* or 202101* or 202102*).dc. use oomezd [time limits in Embase]</p> <p>8 (1 and (3 or 5) and 6) use medall</p> <p>9 (2 and (4 or 5) and 7) use oomezd</p>
--	--	---

### 6.3 Search strategy to identify ongoing studies


NIPN is responsible for searching in trial registries to identify ongoing and unpublished studies. The combination of search terms related to COVID-19 and tocilizumab are described in Appendix Table 6-3.

**Table 6-3 Search strategy to identify ongoing studies**

Database	URL	Search line / search terms	Date of search	Hits retrieved
ClinicalTrials.gov	<a href="https://clinicaltrials.gov/">https://clinicaltrials.gov/</a>	<p>"Basic search mode"</p> <p>Terms used at Condition or disease:</p> <ul style="list-style-type: none"> <li>• covid19</li> </ul> <p>Terms used at "other terms":</p> <ul style="list-style-type: none"> <li>• tocilizumab</li> <li>• Actemra</li> </ul>	09/03/2021	54 2 new
ISRCTN	<a href="https://www.isrctn.com/">https://www.isrctn.com/</a>	<p>Basic search mode</p> <p>Search terms:</p> <ol style="list-style-type: none"> <li>1. covid-19 and tocilizumab</li> <li>2. covid-19 and Actemra</li> <li>3. SARS-CoV-2 and tocilizumab</li> <li>4. SARS-CoV-2 and Actemra</li> </ol>	09/03/2021	2 0 new
European Clinical Trials Registry	<a href="https://www.clinicaltrialsregister.eu/">https://www.clinicaltrialsregister.eu/</a>	<p>Basic search mode</p> <p>Search terms:</p> <ol style="list-style-type: none"> <li>1. covid-19 and tocilizumab</li> <li>2. covid-19 and Actemra</li> <li>3. SARS-CoV-2 and tocilizumab</li> <li>4. SARS-CoV-2 and Actemra</li> </ol>	09/03//2021	33 0 new

\* In Basic Search mode, one term was added to the field "condition or disease" and one term in the field "other terms".


## 6.4 Flow diagrams



### Appendix Figure 6-1. Flow diagram depicting the selection process of RCTs

RCT = randomised controlled trial;

\* The selection process was part of an external project, see <https://www.deplazio.net/farmacicovid> and Prospero ID CRD42020176914.



**Appendix Figure 6-2. Flow diagram depicting the selection process of observational studies**  
\*\* studies evaluating active substances relevant to other EUnetHTA rolling collaborative reviews